

**2005Table 8-5. Hypersensitivity pneumonitis: Number of deaths, mortality rates (per million population), and years of potential life lost (YPLL) by state, U.S. residents age 15 and over, 1993–2002**

State	No. of Deaths		Crude Death Rate		Age-Adjusted Death Rate		YPLL to Life Expectancy			
	Deaths	Rank	Rate	Rank	Rate	Rank	Total	Rank	YPLL/death	Rank
Alabama	3	36	0.09	41	0.09	42	45	36	15.1	21
Alaska	–	–	–	–	–	–	–	–	–	–
Arizona	4	27	0.11	38	0.11	37	105	20	26.4	1
Arkansas	2	39	0.10	39	0.09	42	36	39	17.9	12
California	53	1	0.21	21	0.23	17	935	1	17.6	14
Colorado	10	14	0.32	14	0.36	14	179	13	17.9	12
Connecticut	6	21	0.23	18	0.22	19	127	17	21.2	5
Delaware	1	44	0.17	29	0.18	28	21	43	21.1	6
District of Columbia	1	44	0.23	18	0.23	17	9	47	8.6	46
Florida	23	7	0.19	26	0.15	30	317	7	13.8	29
Georgia	2	39	0.03	48	0.05	47	17	44	8.5	47
Hawaii	–	–	–	–	–	–	–	–	–	–
Idaho	8	18	0.85	4	0.91	3	97	21	12.1	36
Illinois	17	10	0.18	28	0.18	28	245	11	14.4	24
Indiana	7	20	0.15	31	0.15	30	93	23	13.3	31
Iowa	24	5	1.05	2	0.89	5	268	9	11.2	43
Kansas	9	16	0.44	12	0.43	12	170	14	18.9	8
Kentucky	4	27	0.13	33	0.13	33	87	25	21.7	2
Louisiana	4	27	0.12	36	0.12	34	74	29	18.5	10
Maine	4	27	0.40	13	0.37	13	46	34	11.5	39
Maryland	4	27	0.10	39	0.11	37	53	32	13.2	32
Massachusetts	4	27	0.08	44	0.08	44	46	34	11.5	39
Michigan	22	9	0.29	15	0.30	15	334	6	15.2	19
Minnesota	23	7	0.62	9	0.61	9	281	8	12.2	35
Mississippi	1	44	0.05	47	0.05	47	14	45	14.4	24
Missouri	9	16	0.21	21	0.20	25	151	15	16.7	15
Montana	2	39	0.29	15	0.28	16	23	42	11.3	41
Nebraska	8	18	0.62	9	0.56	10	97	21	12.1	36
Nevada	–	–	–	–	–	–	–	–	–	–
New Hampshire	2	39	0.21	21	0.22	19	28	41	14.2	28
New Jersey	5	24	0.08	44	0.08	44	78	28	15.5	18
New Mexico	1	44	0.08	44	0.08	44	14	45	14.4	24
New York	32	3	0.22	20	0.22	19	410	3	12.8	33
North Carolina	12	12	0.20	24	0.21	23	136	16	11.3	41
North Dakota	4	27	0.79	5	0.68	7	74	29	18.6	9
Ohio	17	10	0.19	26	0.19	27	268	9	15.8	16
Oklahoma	4	27	0.15	31	0.15	30	60	31	14.9	23
Oregon	12	12	0.46	11	0.44	11	126	18	10.5	44
Pennsylvania	24	5	0.25	17	0.22	19	362	5	15.1	21
Rhode Island	1	44	0.13	33	0.10	39	8	48	8.3	48
South Carolina	6	21	0.20	24	0.21	23	86	26	14.4	24
South Dakota	6	21	1.04	3	0.94	2	82	27	13.6	30
Tennessee	5	24	0.12	36	0.12	34	107	19	21.4	4
Texas	26	4	0.17	29	0.20	25	395	4	15.2	19
Utah	10	14	0.66	7	0.83	6	194	12	19.4	7
Vermont	3	36	0.63	8	0.67	8	38	38	12.5	34
Virginia	5	24	0.09	41	0.10	39	91	24	18.2	11
Washington	4	27	0.09	41	0.10	39	47	33	11.8	38
West Virginia	2	39	0.13	33	0.12	34	43	37	21.6	3
Wisconsin	45	2	1.09	1	1.07	1	705	2	15.7	17
Wyoming	3	36	0.79	5	0.90	4	29	40	9.7	45

– indicates no deaths listed.

NOTE: Slight variations in numbers may exist when compared to previous WoRLD Surveillance Reports. See appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Bureau of the Census.